

# Euclid's Algorithm for the Greatest Common Divisor

Desh Ranjan

Department of Computer Science

New Mexico State University

## 1 Numbers, Division and Euclid

People have been using numbers, and operations on them like division, for a very long time for practical purposes like dividing up the money left by parents for children, or distributing ears of corn equally to groups of people, and more generally to conduct all sorts of business dealings. It may be a bit of a surprise that things like calculating divisors of numbers also form the core of today's methods ensuring security of computer systems and internet communications. The RSA cryptosystem that is used extensively for secure communications is based on the assumed difficulty of calculating divisors of large numbers, so calculating divisors is important even today.

A related and even more basic notion is that of multiples of quantities. A natural way to compare quantities is to "measure" how many times we need to aggregate the smaller quantity to obtain the larger quantity. For example, we may be able to compare two unknown lengths by observing that the larger length can be obtained by "aggregating" the smaller length three times. This provides a sense of how the two lengths compare without actually knowing the two lengths.

The larger quantity may not always be obtainable from the smaller quantity by aggregating it an integral number of times. In this scenario, one way to think would be to imagine each of the two quantities to be made up of smaller (identical) parts such that both the quantities can be obtained by aggregating these smaller parts an integral number of times. Obviously, we will need a greater number of these parts for the larger quantity than for the smaller one. For example, when comparing two weights, one might observe that the larger one can be obtained by aggregating some weight 7 times whereas the smaller weight can be obtained by aggregating the same weight 5 times. This provides a basis for comparing the two weights. Of course, in the above scenario, one can also observe that if we chose even smaller parts to "split" the weights (say a quarter of the first one), the first weight would be obtained by aggregating this even smaller weight 28 times and the smaller of the two original weights would be obtained by aggregating this smaller part 20 times, which also provides us a sense of the relative magnitudes of

the two weights. However, using smaller numbers like 7 and 5 to describe relative magnitudes seems intuitively and practically more appealing than using larger numbers, like 28 and 20. This leads us to think about what would be the greatest magnitude such that two given magnitudes will both be multiples of that common magnitude.

This question was considered by Greek mathematicians more than 2000 years ago. One of those Greeks was Euclid, who compiled a collection of mathematical works called *Elements* that has a chapter, interestingly called a “Book”, about numbers. During the course of this project you will read a translation of part of this chapter to discover Euclid’s method (algorithm) to compute the greatest common divisor of two numbers. It is not clear if Euclid was the first person to discover this algorithm, but his is the earliest known written record of it.

## 1.1 Euclid of Alexandria

Euclid lived around 300 B.C.E. Very little is known about his life. It is generally believed that he was educated under students of Plato’s Academy in Athens. According to Proclus (410–485 C.E.), Euclid came after the first pupils of Plato and lived during the reign of Ptolemy I (306–283 B.C.E.). It is said that Euclid established a mathematical school in Alexandria. Euclid is best known for his mathematical compilation *Elements* in which among other things he laid down the foundations of geometry and number theory. The geometry that we learn in school today traces its roots to this book, and Euclid is sometimes called the father of geometry.

Euclid did not study mathematics for its potential practical applications or financial gains. He studied mathematics for a sense of order, structure and the ideal form of reason. To him geometrical objects and numbers were abstract entities, and he was interested in studying and discovering their properties. In that sense, he studied mathematics for its own sake. One story that reveals his disdain for learning for the purpose of material gains concerns a pupil who had just finished his first geometry lesson. The pupil asked what he would gain from learning geometry. As the story goes, Euclid asked his subordinate to give the pupil a coin so that he would be gaining from his studies. Another story that reveals something about his character concerns King Ptolemy. Ptolemy asked the mathematician if there was an easier way to learn geometry. Euclid replied, “There is no royal road to geometry”, and sent the king to study.

Euclid wrote several books such as *Data*, *On Divisions of Figures*, *Phaenomena*, *Optics*, and the lost books *Conics* and *Porisms*, but *Elements* remains

his best known compilation. The first “book” [chapter] in this compilation is perhaps the most well-known. It lays down the foundations of what we today call “Euclidean” geometry (which was the only plane geometry people studied until the Renaissance). This book has definitions of basic geometric objects like points and lines along with basic postulates or axioms. These axioms are then used by Euclid to establish many other truths (*Theorems*) of geometry. Euclid’s *Elements* is considered one of the greatest works of mathematics, partly because it is the earliest we have that embodies an axiomatic approach. It was translated into Latin and Arabic and influenced mathematics throughout Europe and the Middle East. It was probably the standard “textbook” for geometry for more than 1500 years in western Europe and continues to influence the way geometry is taught to this day.

Book 7 of *Elements* provides foundations for number theory. Euclid’s Algorithm for calculating the greatest common divisor of two numbers was presented in this book. As one will notice later, Euclid uses lines to represent numbers and often relies on visual figures to aid the explanation of his method of computing the greatest common divisor (GCD) of two numbers. As such, he seems to be relating numbers to geometry, which is quite different from the present day treatment of number theory.

Today, erroneously, many different methods are called Euclid’s algorithm. By reading the original writings of Euclid you will discover the real Euclidean algorithm and appreciate its subtlety. In any case, “Euclid’s Algorithm” is one of the most cited and well-known examples of an (early) algorithm. To quote Knuth [2] :

*By 1950, the word algorithm was mostly associated with “Euclid’s Algorithm”.*

## 2 Prelude

We say that a number<sup>1</sup>  $x$  divides another number  $y$  if  $y$  is a multiple of  $x$ . For example, 1, 2, and 3 all divide 6 but 5 does not divide 6. The only divisors of 17 are 1 and 17. The notation  $x|y$  is a shorthand for “ $x$  divides  $y$ ”. We denote by  $divisors(x)$  the set of all the numbers  $y$  such that  $y|x$ . So, for example,  $divisors(6) = \{1, 2, 3, 6\}$  and  $divisors(60) = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$ .

A number  $z$  is called a common divisor of two numbers  $x$  and  $y$  if  $z|x$  and  $z|y$ . We denote by  $cd(x, y)$  the set of all common divisors of  $x$  and  $y$ .

---

<sup>1</sup>The word number in this section means a positive integer. That is what it meant to Euclid.

For example,  $cd(6, 8) = \{1, 2\}$  and  $cd(40, 180) = \{1, 2, 4, 5, 10, 20\}$ .

**Exercise 2.1.** What is the set of divisors of the number 315?

**Exercise 2.2.** Calculate the set  $cd(288, 216)$ .

While it is relatively easy to calculate the divisors of a number and common divisors of two numbers when the numbers are small, the task becomes harder as the numbers becomes larger.

**Exercise 2.3.** Calculate  $divisors(3456)$ .

**Exercise 2.4.** Calculate  $cd(3456, 4563)$ .

**Exercise 2.5.** A rather naive method for computing the divisors of a number  $x$  is to test whether each number from 1 to  $x$  inclusive is a divisor of  $x$ . For integers  $n = 1, 2, 3, \dots, x$ , simply test whether  $n$  divides  $x$ . Using this naive algorithm, write a computer program in the language of your choice that accepts as input a positive integer  $x$  and outputs all divisors of  $x$ . Run this program for:

- (a)  $x = 3456$ ,
- (b)  $x = 1009$ ,
- (c)  $x = 1080$ .


**Exercise 2.6.** The naive method for computing the common divisors of two numbers  $x$  and  $y$  is to test whether each number from 1 to the least of  $\{x, y\}$  divides  $x$  and  $y$ . In modern notation, let  $m$  denote the minimum (least of)  $\{x, y\}$ . For  $n = 1, 2, 3, \dots, m$ , first test whether  $n$  divides  $x$ , and, if so, then test whether  $n$  divides  $y$ . If  $n$  divides both  $x$  and  $y$ , record  $n$  as a common divisor. Using this naive algorithm, write a computer program in the language of your choice that accepts as input two positive integers  $x, y$ , and outputs their common divisors. Run this program for:

- (a)  $x = 3456, y = 4563$ ,
- (b)  $x = 625, y = 288$ ,
- (c)  $x = 216, y = 288$ ,
- (d)  $x = 147, y = 27$ .

As you might have noticed the number 1 divides every number. Since there is no number smaller than 1, 1 is the **smallest** common divisor for any two numbers  $x$  and  $y$ . What about the **greatest** common divisor? The greatest common divisor of two numbers  $x$  and  $y$ , denoted by  $gcd(x, y)$ , is the largest number  $z$  such that  $z|x$  and  $z|y$ . Finding the greatest common divisor is not nearly as easy as finding the smallest common divisor.

### 3 Euclid's Algorithm

Here we present the translations of (relevant) Definitions, Proposition 1 and Proposition 2 from Book VII of Euclid's *Elements* as translated by Sir Thomas L. Heath [1]. Euclid's method of computing the GCD is based on these propositions.



#### BOOK VII of *Elements* by Euclid


##### DEFINITIONS.

1. A **unit** is that by virtue of which each of the things that exist is called one.
2. A **number** is a multitude composed of units.
3. A number is a **part** of a number, the less of the greater, when it measures the greater.
4. but **parts** when it does not measure it.<sup>2</sup>
5. The greater number is a **multiple** of the less when it is measured by the less.
6. An **even number** is that which is divisible into two equal parts.
7. An **odd number** is that which is not divisible into two equal parts, or that differs by a unit from an even number.
8. An **even-times even number** is that which is measured by an even number according to an even number.

---

<sup>2</sup>While this definition is not relevant here, what is meant by this definition is quite subtle and the subject of scholarly mathematical work.

9. An **even-times odd number** is that which is measured by an even number according to an odd number.
10. An **odd-times odd number** is that which is measured by an odd number according to an odd number.
11. A **prime number** is that which is measured by a unit alone.<sup>3</sup>
12. Numbers **prime to one another** are those which are measured by a unit alone as a common measure.
13. A **composite number** is that which is measured by some number.
14. Numbers **composite to one another** are those which are measured by some number as a common measure.



**Exercise 3.1.** Discuss how Euclid’s “unit” relates to the number 1. Does Euclid think that 1 is a number?

**Exercise 3.2.** What is likely meant when Euclid states that a number “measures” another number? Express Euclid’s notion of “measures” in modern mathematical notation.

**Exercise 3.3.** Does the number 4 measure number 72? Does 5 measure 72? Briefly justify your answer.

**Exercise 3.4.** Euclid never defines what is a “common measure,” but uses that in definition 12 and 14. What is your interpretation of Euclid’s “common measure”?


**Exercise 3.5.** Find a number (other than the unit) that is a common measure of the numbers 102 and 187. According to Euclid’s definitions, are the numbers 102 and 187 composite to one another? Why or why not?

**Exercise 3.6.** According to Euclid’s definitions, are the numbers 21 and 55 composite to one another? Justify your answer.

We now present Proposition 1 from Euclid’s book VII. The proposition concerns numbers that are prime to one another.

---

<sup>3</sup>Reading further work of Euclid, e.g. Proposition 2, it is clear that Euclid meant that a prime number is that which is measured only by the unit and the number itself.




PROPOSITION 1.

*Two unequal numbers being set out, and the less being continually subtracted in turn from the greater, if the number which is left never measures the one before it until a unit is left, the original numbers will be prime to one another.*

For, the less of two unequal numbers  $AB$ ,  $CD$  being continually subtracted from the greater, let the number which is left never measure the one before it until a unit is left;

I say that  $AB$ ,  $CD$  are prime to one another, that is, that a unit alone measures  $AB$ ,  $CD$ .

For, if  $AB$ ,  $CD$  are not prime to one another, some number will measure them.



Let a number measure them, and let it be  $E$ ; let  $CD$ , measuring  $BI$ , leave  $IA$  less than itself,

let,  $AI$  measuring  $DG$ , leave  $GC$  less than itself,

and let  $GC$ , measuring  $IH$ , leave a unit  $HA$ .

Since, then  $E$  measures  $CD$ , and  $CD$  measure  $BI$ , therefore  $E$  also measures  $BI$ .

But it also measures the whole  $BA$ ;

therefore it will also measure the remainder  $AI$ .

But  $AI$  measures  $DG$ ;

therefore  $E$  also measures  $DG$ .

But it also measures the whole  $DC$ ;

therefore it will also measure the remainder  $CG$ .

But  $CG$  measures  $IH$ ;


therefore  $E$  also measures  $IH$ .

But it also measures the whole  $IA$ ;

therefore it will also measure the remainder, the unit  $AH$ , though it is a number: which is impossible.

Therefore no number will measure the numbers  $AB$ ,  $CD$ ; therefore  $AB$ ,  $CD$  are prime to one another. [VII. Def 12]

Q. E. D.



**Exercise 3.7.** Euclid begins with two unequal numbers  $AB$ ,  $CD$ , and continually subtracts the smaller in turn from the greater. Let's examine how this method proceeds "in turn" when subtraction yields a new number that is smaller than the one subtracted. Begin with  $AB = 162$  and  $CD = 31$ .

- (a) How many times must  $CD$  be subtracted from  $AB$  until a remainder is left that is less than  $CD$ ? Let this remainder be denoted as  $IA$ .
- (b) Write  $AB = BI + IA$  numerically using the given value for  $AB$  and the computed value for  $IA$ .
- (c) How many times must  $IA$  be subtracted from  $CD$  until a remainder is left that is less than  $IA$ ? Let this remainder be denoted as  $GC$ .



- (d) Write  $CD = DG + GC$  numerically using the given value for  $CD$  and the computed value for  $GC$ .
- (e) How many times must  $GC$  be subtracted from  $IA$  until a remainder is left that is less than  $GC$ ? Let this remainder be denoted as  $HA$ .
- (f) Is  $HA$  a unit?
- (g) Write  $IA = IH + HA$  numerically using the computed values of  $IA$  and  $HA$ .


**Exercise 3.8.** Apply the procedure outlined in Proposition 1 to the numbers  $AB = 625$  and  $CD = 288$ . Begin by answering questions (a)–(f) above except with the new values for  $AB$  and  $CD$ .

- (g) In this example, how should the algorithm proceed until a remainder is reached that is a unit?

**Exercise 3.9.** Euclid claims that if the repeated subtraction algorithm of Proposition 1 eventually produces a unit as a remainder, then the original numbers  $AB, CD$  are prime to one another. He does so by using a “proof by contradiction.” Suppose the result, namely that  $AB$  and  $CD$  are prime to one another, is false. In this exercise we examine the consequences of this.

- (a) If  $AB$  and  $CD$  are not prime to one another, must these numbers have a common measure  $E$  that is greater than 1? Justify your answer by using Euclid’s definitions.
- (b) From  $AB = BI + IA$ , why must  $E$  also measure  $IA$ ? Be sure to carefully justify your answer for general numbers  $AB$  and  $CD$  (not tied to one particular example).
- (c) From  $CD = DG + GC$ , why must  $E$  also measure  $GC$ ? Be sure to carefully justify your answer.
- (d) From  $IA = IH + HA$ , why must  $E$  also measure  $HA$ ? Carefully justify your answer.
- (e) If according to Euclid,  $HA$  is a unit, what contradiction has been reached in part (d)?

We now present proposition 2 from Book VII of Euclid's elements. This proposition presents a method to compute the GCD of two numbers which are not prime to each other and provides a proof of the correctness of the method. Euclid's presentation intermixes the proof and the method to some extent. Despite this the elegance of his method and the proof is striking.



### PROPOSITION 2.

*Given two numbers not prime to one another, to find their greatest common measure.*

Let  $AB$ ,  $CD$  be the two given numbers not prime to one another.

Thus it is required to find the greatest common measure of  $AB$ ,  $CD$ .

If now  $CD$  measures  $AB$  - and it also measures itself -  $CD$  is a common measure of  $CD$ ,  $AB$ .

And it is manifest that it is also the greatest; for no greater number than  $CD$  will measure  $CD$ .

But, if  $CD$  does not measure  $AB$ , then, the less of the numbers  $AB$ ,  $CD$  being continually subtracted from the greater, some number will be left which will measure the one before it.<sup>4</sup>

For a unit will not be left; otherwise  $AB$ ,  $CD$  will be prime to one another [VII, I], which is contrary to the hypothesis.


Therefore some number will be left which will measure the one before it.

Now let  $CD$ , measuring  $BE$ , leave  $EA$  less than itself, let  $EA$ , measuring  $DI$ , leave  $IC$  less than itself, and let  $CI$  measure  $AE$ .

Since then,  $CI$  measures  $AE$ , and  $AE$  measures  $DI$ ,

---

<sup>4</sup>This is the heart of Euclid's description of his algorithm. The statement is somewhat ambiguous and subject to at least two different interpretations.



therefore  $CI$  will also measure  $DI$ .

But it also measures itself;

therefore it will also measure the whole  $CD$ .

But  $CD$  measures  $BE$ ;

therefore  $CI$  also measures  $BE$ .

But it also measures  $EA$ ;

therefore, it will also measure the whole  $BA$ .

But it also measures  $CD$ ;

therefore  $CI$  measures  $AB, CD$ .

Therefore  $CI$  is a common measure of  $AB, CD$ .

I say next that it is also the greatest.

For, if  $CI$  is not the greatest common measure of  $AB, CD$ , some number which is greater than  $CI$  will measure the numbers  $AB, CD$ .

Let such a number measure them, and let it be  $G$ .

Now, since  $G$  measures  $CD$ , while  $CD$  measures  $BE$ ,  $G$  also measures  $BE$ .

But it also measures the whole  $BA$ ;

therefore it will also measure the remainder  $AE$ .

But  $AE$  measures  $DI$ ;

therefore  $G$  will also measure  $DI$ .


But it will also measure the whole  $DC$ ;

therefore it will also measure the remainder  $CI$ , that is, the greater will measure the less: which is impossible.

Therefore no number which is greater than  $CI$  will measure the numbers  $AB$ ,  $CD$ ;

therefore  $CI$  is the greatest common measure of  $AB$ ,  $CD$ .

**PORISM.** From this it is manifest that, if a number measure two numbers, it will also measure their greatest common measure.



**Exercise 3.10.** In Proposition 2 Euclid describes a procedure to compute the greatest common measure of two numbers  $AB$ ,  $CD$ , not prime to one another. The method again proceeds by repeatedly subtracting the smaller in turn from the greater until some number is left, which in this case divides the number before it. Let's examine this process for  $AB = 147$  and  $CD = 27$ .

- (a) Does  $CD$  measure  $AB$ ? If so, the process stops. If not, how many times must  $CD$  be subtracted from  $AB$  until a positive remainder is left that is less than  $CD$ . Let  $EA$  denote this remainder.
- (b) Write  $AB = BE + EA$  numerically using the given value for  $AB$  and the computed value for  $EA$ . Also find a positive integer  $q_1$  so that  $BE = q_1 \cdot CD$ .

- (c) Does  $EA$  measure  $CD$ ? If so, the process stops. If not, how many times must  $EA$  be subtracted from  $CD$  until a positive remainder is left that is less than  $EA$ . Let  $IC$  denote this remainder.
- (d) Write  $CD = DI + IC$  numerically using the given value for  $CD$  and the computed value for  $IC$ . Also, find a positive integer  $q_2$  so that  $DI = q_2 \cdot EA$ .
- (e) Does  $IC$  measure  $EA$ ? If so, the process stops. If not, how many times must  $IC$  be subtracted from  $EA$  until a positive remainder is left that is less than  $IC$ ?
- (f) Find a positive integer  $q_3$  so that  $EA = q_3 \cdot IC$ .

**Exercise 3.11.** Apply Euclid's procedure in Proposition 2 to compute the greatest common measure of  $AB = 600$  and  $CD = 276$  outlined in the steps below.

- (a) To streamline the process, let  $a_1 = AB = 600$ ,  $a_2 = CD = 276$ , and  $a_3 = EA$ . Compute  $a_3$  numerically for this example. Write the equation  $AB = BE + EA$  entirely in terms of  $a_1$ ,  $a_2$  and  $a_3$ .
- (b) Let  $a_4 = IC$ . Compute  $a_4$  for this example. Write the equation  $CD = DI + IC$  entirely in terms of  $a_2$ ,  $a_3$  and  $a_4$ .
- (c) Does  $IC$  measure  $EA$  in this example? If so, the process stops. If not, how many times must  $IC$  be subtracted from  $EA$  until a positive remainder is left that is less than  $IC$ ? Denote this remainder by  $a_5$ .
- (d) Write an equation using  $a_3$ ,  $a_4$  and  $a_5$  that reflects the number of times  $IC$  must be subtracted from  $EA$  so that the remainder is  $a_5$ .
- (e) Does  $a_5$  measure  $a_4$ ? If so, the process stops. If not, how many times must  $a_5$  be subtracted from  $a_4$  until a positive remainder is left that is less than  $a_5$ ?

**Exercise 3.12.** In modern notation, the Euclidean algorithm to compute the greatest common measure of two positive integers  $a_1$  and  $a_2$  (prime to each other or not) can be written as follows. Find a sequence of positive integer remainders  $a_3, a_4, a_5, \dots, a_{n+1}$  and a sequence of (positive) integer

multipliers  $q_1, q_2, q_3, \dots, q_n$  so that

$$\begin{aligned}
 a_1 &= q_1 a_2 + a_3, & 0 < a_3 < a_2 \\
 a_2 &= q_2 a_3 + a_4, & 0 < a_4 < a_3 \\
 a_3 &= q_3 a_4 + a_5, & 0 < a_5 < a_4 \\
 &\vdots \\
 a_{i-1} &= q_{i-1} a_i + a_{i+1}, & 0 < a_{i+1} < a_i \\
 a_i &= q_i a_{i+1} + a_{i+2}, & 0 < a_{i+2} < a_{i+1} \\
 &\vdots \\
 a_{n-1} &= q_{n-1} a_n + a_{n+1}, & 0 < a_{n+1} < a_n \\
 a_n &= q_n a_{n+1}
 \end{aligned}$$

- (a) Why is  $a_{n+1}$  a divisor of  $a_n$ ? Briefly justify your answer.
- (b) Why is  $a_{n+1}$  a divisor of  $a_{n-1}$ ? Carefully justify your answer.
- (c) In a step-by-step argument, use (backwards) mathematical induction to verify that  $a_{n+1}$  is a divisor of  $a_i$ ,  $i = n, n-1, n-2, \dots, 3, 2, 1$ .
- (d) Why is  $a_{n+1}$  a common divisor of  $a_1$  and  $a_2$ ?
- (e) In a step-by-step argument, use (forwards) mathematical induction to verify that if  $G$  is a divisor of  $a_1$  and  $a_2$ , then  $G$  is also a divisor of  $a_i$ ,  $i = 3, 4, 5, \dots, n+1$ . First, carefully explain why  $G$  is a divisor of  $a_3$ . Then examine the inductive step.
- (f) From part (d) we know that  $a_{n+1}$  is a common divisor of  $a_1$  and  $a_2$ . Carefully explain how part (e) can be used to conclude that  $a_{n+1}$  is in fact the *greatest* common divisor of  $a_1$  and  $a_2$ . A proof by contradiction might be appropriate here, following Euclid's example.

**Exercise 3.13.** In Proposition 1 Euclid describes an algorithm whereby, given two unequal numbers, the less is continually subtracted in turn from the greater until a unit is left. While in Proposition 2, Euclid describes an algorithm, whereby, given two unequal numbers, the less is continually subtracted from the greater until some number is left which measures the one before it.

- (a) To what extent are these algorithms identical?

- (b) How are the algorithms in Proposition 1 and Proposition 2 designed to differ in application?
- (c) Does Euclid consider a unit as a number? Justify your answer citing relevant passages from the work of Euclid. Does Euclid consider a common measure as a number? Again, justify your answer from the work of Euclid.
- (d) Why, in your opinion, does Euclid describe this algorithm using two separate propositions, when a single description could suffice?

**Exercise 3.14.** In the modern description of the Euclidean algorithm in Exercise (3.12), the last equation written is

$$a_n = q_n a_{n+1},$$

meaning that after  $n$ -steps, the algorithm halts and  $a_{n+1}$  divides (measures)  $a_n$ . Given any two positive integers  $a_1$  and  $a_2$ , why must the Euclidean algorithm halt in a finite number of steps? Carefully justify your answer using the modern version of the algorithm.

**Exercise 3.15.** Write a computer program in the language of your choice that implements Euclid's algorithm for finding the greatest common divisor of two positive integers. The program should accept as input two positive integers  $a_1, a_2$ , and as output print their greatest common divisor. Run the program for:

- (a)  $a_1 = 3456, a_2 = 4563$ ,
- (b)  $a_1 = 625, a_2 = 288$ ,
- (c)  $a_1 = 216, a_2 = 288$ .

## 4 Notes to the Instructor

The project is meant for use for in an introductory computer science or discrete mathematics class. The project can be used to introduce students to the notion of “computation method” or “algorithm and to explore concepts like iteration in a basic setting. It allows them to practice their skills in doing proofs but more importantly to observe the evolution of what is accepted as a valid proof or a well-described algorithm. The students will easily notice that the method presented by Euclid to compute the GCD and proof of

its correctness that he provided would not be formally accepted as correct today. They will also notice, however, that Euclid is somehow able to convey his ideas behind his method and proof in a way that they can be easily translated into a modern algorithm and proof of its correctness. In this way, it will provide them a sense of connection to the past.

A basic knowledge of programming is essential to successfully complete some of the components of the project.

## References

- [1] Heath, T., *Euclid: The Thirteen Books of the Elements*, Volume 2, Second Edition, Dover Publications, New York, 1956.
- [2] Knuth, D., *The Art of Computer Programming*, Volume 1, Addison-Wesley, Reading, Mass., 1968.